

Rx.



Stillen[®] Tab.
[Eupatilin]

A Novel Drug for Prevention of
NSAIDs-induced Gastritis

 **DONG-A ST**

Characteristics of Stillen®


- A Novel Cytoprotectant for Gastritis
- Triple Modes of Action
- Indicated for Prevention of NSAIDs-induced Gastritis

Indication of Stillen®



Excellent Efficacy on Erosive Gastritis

Stillen® shows significantly higher improvement rates & cure rates of erosive gastritis compared to cetraxate among the patients [Phase III]¹⁾




* An active-controlled, double-blinded, non-inferiority trial conducted in South Korea
 Ref 1) S.Y. Seol et al. World J Gastroenterol 2004 Aug 15;10(16):2379-82

A Novel Drug For Prevention of NSAIDs-induced Gastritis



Proven Efficacy on Prevention of NSAIDs-induced Gastritis


Stillen® shows non-inferiority in gastric mucosa protection rates and no difference in duodenal mucosa protection rates compared to misoprostol among the patients. [Phase IV]²⁾



* A multicenter, double-blinded, stratified randomized, active-controlled, parallel group, non-inferiority trial conducted in South Korea from Sep. 2012 to Mar. 2014

Ref 2) O. Y. Lee et al. Archives of Pharmacal Research, 2014 Oct; 37(10):1308-1316

Stillen® shows non-inferiority in gastric mucosa protection rates and no difference in duodenal mucosa protection rates compared to misoprostol among the healthy subjects [Phase IV]³⁾



* A multicenter, double-blinded, randomized non-inferiority trial conducted in South Korea.


Ref 3) K. N. Lee et al. J. Korean Med Sci. 2011 Aug;26(8): 1074-80

Stillen[®] Tab.



Favorable Safety Profile

Stillen[®] has more favorable safety profile than misoprostol in the incidence rate of GI-related adverse events[Phase IV]⁴⁾



* A randomized, multicenter, double-blinded study comparing Stillen[®] with misoprostol
 Ref 4) O. Y. Lee et al. Archives of Pharmacal Research, 2014 Oct; 37(10):1308-1316

Composition

For each tablet (339.25mg) contains
 95% ethanol soft extracts (20 → 1) of Artemisiae Argyi Folium 60mg
 (0.48~1.44mg as Eupatillin and 0.15~0.45mg as Jaceosidin)

Description

Green, oval-shaped, film coated tablet.

Indications

Improvement of gastric mucosal lesion(erosion, hemorrhage, reddening, edema)
 in acute and chronic gastritis, prevention of NSAIDs-induced gastritis

Dosage and administration

Take one tablet three times a day

Storage

Air-tight container at room temperature(1~30°C)



DONG-A ST

64, Cheonho-daero, Dongdaemun-gu, Seoul, South Korea
 Tel: +82-2-920-8399 Fax: +82-2-924-2662 <http://en.donga-st.com>

SLT-1509-OS Designed by Overseas PMA Team Printed by TAMA